

COOLING A HOT DISK AROUND A SUPERMASSIVE BLACK HOLE BY A STAR

Heon-Young Chang

Korea Institute For Advanced Study, Seoul, 130-012, Korea

`hyc@kias.re.kr`

Received _____; accepted _____

ABSTRACT

If a supermassive black hole resides in the centers of galaxies, there are several effects expected to be observed. It is likely that accretion disks are around the supermassive black hole. Stellar interactions with the accretion flows around the supermassive black hole play a role in that an flying-by star may cool a hot accretion disk as a result of Comptonization. It is shown that the Comptonization of the stellar emission will take place in a hot accretion disk such as the ADAFs around the supermassive black hole and become a relatively important source of the accretion disk cooling when the mass accretion rate is low, and stars are passing outer parts of the hot accretion disk. We suggest that such a stellar cooling can be observed in the radio frequency regime since synchrotron luminosity of the ADAFs depends strongly on the electron temperature and occurs much more frequently than a tidal disruption event.

Subject headings: accretion, accretion disk – black hole physics

1. INTRODUCTION

Since the early days of research on quasars and AGNs, supermassive black holes (SMBHs) have been considered as the most likely power sources of the activity in these objects (Lynden-Bell 1969; Rees 1984). Many lines of direct and indirect evidence suggest that SMBHs reside in the centers of a substantial fraction of galaxies (Kormendy & Richstone 1995). If this is the case, several effects are expected to be observed. When stars approach to the SMBH in the center of galaxies, stars can be captured or tidally disrupted by the SMBH in galactic nuclei, resulting in flare-like activity. Several aspects of disruptive stellar encounters with supermassive black holes have been studied both theoretically and

observationally (e.g., Nolthenius & Katz 1982; Bicknell & Gingold 1983; Carter & Luminet 1983; Peterson & Ferland 1986; Hills 1988; Kochanek 1992; 1994; Syer & Clarke 1992; 1993; Laguna et al. 1993; Brandt et al. 1995; Renzini et al. 1995; Storchi-Bergmann et al. 1995; Marck et al. 1996). During such a dramatic event, energy outbursts are expected in different processes and its observational implications are various. Explosive release of nuclear energy due to the extreme compression for the orbiting star may occur (Luminet & Pichon 1989; Luminet & Barbury 1990) and may lead to observable enrichment of heavy elements in the galactic center. Tidal disruption events are rare, but seemingly unavoidable consequences of $10^6 - 10^8 M_\odot$ SMBHs in galactic centers. About half of the stellar debris is bound and accretes onto the central SMBH, and the rest is flung-out. The bound half may create a bright UV/X-ray flare as it accretes onto the SMBHs in a timescale of less than a couple of years (Rees 1988). The detection of a single tidal disruption event could even allow for a crude estimate of the SMBH mass (e.g., Loeb & Ulmer 1997). Khokhlov & Melia (1996) show that interactions between the high velocity unbound material and the interstellar medium (ISM) may result in a supernova-like remnant with properties similar to those of Sagittarius A East. In extremely low angular momentum encounters, the material could deposit over 10^{52} ergs into the ISM, creating a shell like a supernova remnant. More typically, the energy deposited would be $\sim 10^{50}$ ergs. Interaction of orbiting debris may occur after the most tightly bound debris begins its another orbit (Kim et al. 1999).

If there is an accretion disk around the SMBH, a less dramatic process may also occur due to the interactions of a flying-by star and the accretion disk around the SMBH (e.g., Syer et al. 1991; Hall et al. 1996). The process has become interesting particularly when the accretion disk is geometrically thick and hot. The accreted material could be gas expelled from stars via stellar winds and supernovae, or intergalactic gas captured by the galaxy. As the gas accretes, the ions are heated by viscous dissipation and transfer their energy to the electrons so that the energy is radiated. However, the energy is inefficiently transferred

from the ions to the electrons when the mass accretion rate is below a critical rate, and advected inward to the central object (Narayan & Yi 1994). The radiative luminosity of the Advection-Dominated Accretion Flows (ADAFs) is much less than that of the standard cool disk (Shakura & Sunyaev 1973). The ADAFs are optically thin and geometrically thick, i.e., quasi-spherical. The electron temperature is very hot, $\sim 10^9 - 10^{10}$ K and relativistic. Angular velocity of the flow is less than the Keplerian.

In this *Letter* we study the stellar interactions with the ADAFs as a possible cooling mechanism of such a hot accretion disk. This *Letter* begins with descriptions what happens if a star meets a hot accretion disk around the SMBH in § 2. We calculate the heating and cooling rates per volume and present the volume-integrated quantities in § 3. We discuss observational implications and conclude in § 4.

2. WHEN A STAR MEETS A SMBH

Stars can be captured or disrupted by the tidal force of the SMBH in galactic nuclei, depending on the mass of the SMBH, the distance between the star and the SMBH, a ‘compactness’ of the star. Since the tidal radius of the SMBH is proportional to $M_{SMBH}^{1/3}$, while the schwarzschild radius of the SMBH increases proportionally to M_{SMBH} , at a certain mass of the SMBH stars can be swallowed by the SMBH without being disrupted. For a solar type stars this critical mass of the SMBH is about $10^8 M_\odot$. In other words, for $M_{SMBH} \gtrsim 10^8 M_\odot$ a dramatic event, such as the tidal disruption, is expected less often.

What happens if a star passes through the accretion disk around the SMBH? Firstly, dynamical friction causes the viscous heating. The power is given by $P = F_{df} \times v_{rel}$, where F_{df} is the drag force, v_{rel} is the relative velocity of the star with respect to the background gas. Ostriker (1999) has estimated the drag force F_{df} on a star with mass M_* moving

through a uniform gas density ρ with the relative velocity v_{rel} :

$$F_{df} = -4\pi I \left(\frac{GM_*}{v_{rel}} \right)^2 \rho, \quad (1)$$

where the negative sign indicates that the force acts in the opposite direction of the star, G is the gravitational constant. The coefficient I depends on the Mach number, $\mathcal{M} \equiv v_{rel}/c_s$, where c_s is the sound speed of the medium. In the limit of a slow moving $\mathcal{M} \ll 1$, $I_{subsonic} \rightarrow \mathcal{M}^3/3$, so that the resulting F_{df} is proportional to the relative speed of the star. In the limit of a fast moving $\mathcal{M} \gg 1$, $I_{supersonic} \rightarrow \ln(v_{rel}t/r_{min})$, where r_{min} is the effective size of the regime where the gravity of the star dominates. Suppose that the accretion flows around the SMBH occur via the ADAFs. The ADAFs have a quasi-spherical morphology (Narayan & Yi 1995a) and the orbital velocity of the flows is sub-Keplerian. These two effects imply that the magnitude of the hydrodynamic drag is insensitive to the orientation of the stellar motion with respect to the rotation of the accreting gas (cf. Narayan 2000). Under such circumstances, $v_{rel} \sim c_s \sim v_K$, where v_K is the Keplerian speed, and therefore the Mach number is of order unity. We choose the supersonic estimate of I , as it gives an upper limit on the heating due to the drag force. As we set r_{min} to the gravitational capture radius, and $v_{rel}t \rightarrow r_{max}$, where r_{max} is the size of the system, I is given as

$$I \sim \ln \left(\frac{r_{max} v_{rel}^2}{GM_*} \right). \quad (2)$$

By setting $v_{rel} = v_K$, we have

$$I \sim \ln \left(\frac{M_{SMBH}}{M_*} \right). \quad (3)$$

On the other hand, the stellar emission may heat or cool a gaseous medium, depending on the ambient environment. In the ADAFs a star and the stellar motion may enhance the cooling by bremsstrahlung and Comptonization processes. The gas density in front of the star may be increased as the motion of the star may compress the gas. According to Stepney & Guilbert (1983), bremsstrahlung cooling rate per volume is increased as the

density increases, since the bremsstrahlung cooling rate is proportional to the square of the density. Comptonization is also possible because the electrons in the ADAFs are relativistic. Radiation emitted by the star is an important source of soft photons. The spectrum of the emitted radiation can be approximated as a blackbody. The outgoing flux at radius R is given by

$$F_*(R) = \frac{L_*}{4\pi R^2}, \quad (4)$$

where L_* is the stellar luminosity. When the star is at a distance from the central SMBH d , the distance from an arbitrary position in the accretion disk to the star R is related with the distance from the position to the SMBH r as given by

$$R^2 = \left| d^2 + r^2 - 2rd \cos \theta \right|, \quad (5)$$

where θ is the angle between two position vectors **Or** and **Od** with respect to the central SMBH. Now we are in a position to calculate the Comptonization of the stellar flux (cf. Narayan & Yi 1995b). We have the stellar cooling rate as

$$q_{*,C}^- = 3 \frac{F_*}{R} \left(\frac{\theta_e}{x_b} \right)^3 \left\{ \frac{\eta_1}{3} \left[\left(\frac{x_{max}}{\theta_e} \right)^3 - \left(\frac{x_c}{\theta_e} \right)^3 \right] - \frac{\eta_2}{3 + \eta_3} \left[\left(\frac{x_{max}}{\theta_e} \right)^{3+\eta_3} - \left(\frac{x_c}{\theta_e} \right)^{3+\eta_3} \right] \right\}, \quad (6)$$

where $\theta_e = kT_e/m_e c^2$, k being the Boltzmann constant, T_e being the electron temperature, m_e being the electron mass, c being the speed of light, $x_b = h\nu_b/m_e c^2$, h being the Plank constant, $\nu_b = 5.61 \times 10^{10} T_*$, $x_{max} = \max(x_b, 3\theta_e)$, x_c is given by the critical frequency ν_c , η'_k s are defined by Dermer et al. (1991).

The cooling rate of the accretion disk due to the stellar light is dependent upon the position of the star in the accretion flows, and its physical properties, such as, the mass accretion rate, the temperature, the density. The stellar cooling rate per volume due to Comptonization becomes relatively important than those due to other processes of accretion disk cooling when the mass accretion rate becomes small and the star is at large distance.

3. STELLAR COOLING OF DISK

We adopt the following dimensionless variables : mass of the SMBH $m = M/M_{\odot}$; radius from the SMBH $r = R/R_g$, where $R_g = 2GM/c^2 = 2.95 \times 10^5 m \text{ cm}$; and mass accretion rate $\dot{m} = \dot{M}/\dot{M}_{\text{Edd}}$, where $\dot{M}_{\text{Edd}} = L_{\text{Edd}}/\eta_{\text{eff}}c^2 = 1.39 \times 10^{18} m \text{ g s}^{-1}$ (the Eddington accretion rate assuming $\eta_{\text{eff}} = 0.1$). We model the accreting gas as a two-temperature plasma assuming that the flows are spherically symmetric. The quantities of interest are the volume-integrated quantities which are obtained by integrating throughout the volume of the flows. As canonical values in a model for the ADAFs parameters are taken to be $r_{\min} = 3$, $r_{\max} = 10^5$, $\alpha = 0.3$, and $\beta = 0.5$ (see, e.g., Narayan and Yi 1995b). We assume that the star in the accretion disk is a large red giant star, and physical parameters of a large red giant star are quoted from Allen (1973).

Provided that the background gas environment is described by the ADAFs model and that stellar cooling rate per volume due to Comptonization $q_{*,C}^-$ is given by Eq. 6, the volume-integrated cooling rate $dQ_{*,C}^-$ over the spherical shell at r can be obtained as shown in Figure 1. We plot $dQ_{*,C}^-$ with other volume-integrated cooling rates as a function of r . The continuous line represents the $dQ_{*,C}^-$, the dotted line and the dashed line represent volume-integrated cooling rate due to synchrotron cooling $dQ_{\text{sync}}^- + dQ_{\text{sync},C}^-$ and bremsstrahlung cooling $dQ_{br}^- + dQ_{br,C}^-$. Note that the Comptonization of soft photons from synchrotron radiation and bremsstrahlung radiation is included. Various cooling terms are computed such that two energy equations for the ions and for the electrons are met (Narayan & Yi 1995b). In calculating the stellar cooling effect we ignored the bremsstrahlung cooling since the density increase ahead of the stellar motion will result in negligible effects as the bremsstrahlung cooling is a local effect compared with the stellar Compton cooling. Dropping out the bremsstrahlung cooling and taking the heating rate of the supersonic case make the total cooling estimate obtained in this analysis as a lower limit

of the stellar cooling. As shown in Figure 1, the stellar cooling effect becomes important as the mass accretion rate is smaller and as the star locates farther from the central SMBH. Comptonization of stellar photons dominates at the position of the star in the ADAFs when the mass accretion rate is small. In Figure 2 we show that the electron temperature in the left column and the radio spectrum of the ADAFs in the right column. We show the case that the star is at $10 r_g$, that is $r_* = 10$, in Figure 2 as an example. Since the dominant effects on the spectrum is due to the inner parts of the ADAFs, the stellar cooling at closer to the SMBH changes the spectrum more significantly. The electron temperature is averaged over the volume of the shell, and the corresponding synchrotron spectrum of the flows is plotted next to the temperature plot. The continuous line and the dotted line indicate the electron temperature and the radio spectrum of the ADAFs with the stellar cooling, and of the ADAFs without the stellar cooling, respectively. As shown in $\dot{m} = 10^{-5}$ case, Comptonization of stellar soft photons cools out to $r \sim 10^3$, and the spectrum is modified enough to be observed. Since the synchrotron radio spectrum is a strong function of the electron temperature (see Mahadevan 1997), the dominant effect is expected to be observed in the radio regime. The suppression of the radio spectrum due to stellar cooling is the greatest at the frequency corresponding to the position where the star cools the accretion disk. It can be understood by the fact that the synchrotron radio emission of the ADAFs at each frequency is closely related to a specific radius. For instance, the emission at higher frequencies originates at smaller radii, or closer to the central supermassive black hole.

4. DISCUSSION AND CONCLUSION

Physical Consequences of flying-by of a star in a hot accretion disk such as the ADAFs are the increase of the total X-ray flux due to Comptonized photon and the decrease of the

electron temperature and subsequently the radio flux of the hot accretion disk. The X-ray flux change due to the stellar Comptonization is inappreciable. On the other hand, the radio flux change due to the change in the electron temperature of the ADAFs is sensitive enough. In Figure 2 we show the radio spectrum of the ADAF model due to synchrotron emission and the suppression of the luminosity due to the stellar Comptonization. Of course, the conclusion depends on changes of the structure of the accretion disk due to the presence of a star and on responses of the star to the strong gravity of the SMBH. A star interacting with a supermassive black hole becomes vulnerable to expansion and compression which may cause oscillating or enhancement of nuclear reaction rate. The emitted stellar flux is an important factor in this analysis. The actual cooling rate could be also higher because the bremsstrahlung cooling may play a role and because the heating may be less efficient than we estimate in this analysis. As the number of surrounding stars and their distribution are roughly known, the total suppression of the radio flux due to the stellar cooling can be obtained. For instance, suppose $N_* \sim 10^6 - 10^7 \text{ pc}^{-3}$ in a galactic nucleus, the number of stars in $r \sim 10^3 \text{ r}_g$ in the accretion disk around $M_{SMBH} \sim 10^8 M_\odot$ is a few. Thus, the total radio flux will be suppressed further by a factor of few. Note, however, that uncertainty in estimating the total observational radio flux could be involved when stars accumulate on orbits between the influence radius r_h and the core radius r_c , and the density of stars in the cusp bound to the black hole may exceed N_* .

Assuming an isotropic distribution of velocities of the radius r_h at which the central SMBH dynamically influences the stellar motion, then the frequency with which a star would pass within a distance r from the SMBH can be approximated by

$$\approx 10^{-3} M_8^{4/3} \left(\frac{N_*}{10^6 \text{ pc}^{-3}} \right) \left(\frac{\sigma}{300 \text{ km s}^{-1}} \right) \left(\frac{r}{r_t} \right) \text{ yr}^{-1}, \quad (7)$$

where M_8 is the SMBH mass in units of $10^8 M_\odot$, σ is the virial velocity of the stars, r_t is the tidal radius (cf. Rees 1988). It could be used to compare the stellar cooling event rate with

the tidal disruption event rate. For instance, a star is tidally disrupted when the star passes the SMBH within the tidal radius. Therefore, the tidal disruption event may occur once at most every $\sim 10^3$ yrs when the star passes the SMBH with $M_{SMBH} = 10^8 M_\odot$ at a distance of $r = r_t$. The stellar cooling event, however, may occur when a star passes the SMBH at a distance farther than $r = r_t$ so that the stellar event rate could be much larger. In the example we present in the *Letter*, $r_t \approx r_g$ for $M_{SMBH} = 10^8 M_\odot$, and the stellar cooling event rate could be larger than the tidal disruption rate by at least a couple of orders as long as the mass accretion rate is small.

Other observational implications of the stellar cooling are as follows. Firstly, as $dQ_{*,C}^-$ depends on the position of the star in the accretion disk the frequency at which the suppression of the radio flux is the greatest varies according to the stellar positions in the accretion disk. If one may monitor the luminosity changes with the observed frequency one may estimate the stellar orbit in the center of galaxies. Secondly, one may distinguish whether there is a cool and thin accretion disk or there is a hot and thick accretion disk in a galaxy. The stellar cooling process is available only when the electrons are relativistic. Therefore, given that such an observation is made, it could be an evidence of the presence of a hot accretion disk in the center of the galaxy.

The author thank I. Yi for useful discussions and C.-S. Choi for helpful comments.

REFERENCES

- Allen, C. W. 1973, in *Astrophysical Quantities*, 3rd Ed. (London: Athlone Press), Ch. 10.
- Bicknell, G. V., & Gingold, R. A. 1983, *ApJ*, 273, 749
- Brandt, W., N. et al. 1995, *MNRAS*, 273, L47
- Carter, B., & Luminet, J. P. 1983, *A&A*, 121, 97
- Dermer, C. D., Liang, E. P., & Canfield, E. 1991, *ApJ*, 369, 410
- Hall, S. M., Clarke, C. J., & Pringle, J. E. 1996, *MNRAS*, 278, 303
- Hills, J. G. 1988, *Nature*, 331, 687
- Khokhlov, A., & Melia, F. 1996, *ApJL*, 457, L61
- Kim, S., Park, M.-G., & Lee, H.-M. 1999, *ApJ*, 519, 647
- Kochanek, C. S. 1992, *ApJ*, 385, 604
- Kochanek, C. S. 1994, *ApJ*, 422, 508
- Kormendy, J., & Richstone, D. 1995, *ARA&A*, 33, 581
- Laguna et al. 1993, *ApJL*, 410, L83
- Loeb, A., & Ulmer, A. 1997, *ApJ*, 489, 573
- Luminet, J. P., & Pichon, B. 1989, *A&A*, 209, 85
- Luminet, J. P., & Barbury, B. 1990, *AJ*, 99, 838
- Lynden-Bell, D. 1969, *Nature*, 223, 690
- Mahadevan, R. 1997, *ApJ*, 477, 585

- Marck, J. A., et al. 1996, A&A, 306, 666
- Narayan, R. 2000, ApJ, 536, 663
- Narayan, R., & Yi, I. 1994, ApJL, 428, L13
- Narayan, R., & Yi, I. 1995a, ApJ, 444, 231
- Narayan, R., & Yi, I. 1995b, ApJ, 452, 710
- Nolthenius, R. A., & Katz, J. I. 1982, ApJ, 263, 377
- Ostriker, E. C. 1999, ApJ, 513, 252
- Peterson, B. M., & Ferland, G. J. 1986, Nature, 34, 345
- Rees, M. 1984, ARA&A, 22, 471
- Rees, M. 1988, Nature, 333, 523
- Renzini et al. 1995, Nature, 378, 39
- Shakura, N. I., & Sunyaev, R. A. 1973, A&A, 24, 337
- Stepney, S., & Guilbert, P. W. 1983, MNRAS, 204, 1269
- Storchi-Bergmann et al. 1995, ApJ, 443, 617
- Syer, D., & Clarke, C. J. 1992, MNRAS, 255, 92
- Syer, D., Clarke, C. J., & Rees, M. J. 1991, MNRAS, 250, 505
- Syer, D., & Clarke, C. J. 1993, MNRAS, 260, 463

Fig. 1.— The volume integrated cooling rate over the spherical shell due to various mechanisms are shown as a function of r in log scales where $\dot{m} = 10^8$. The continuous line represents the $dQ_{*,C}^-$, the dotted line and the dashed line represent due to synchrotron cooling $dQ_{sync}^- + dQ_{sync,C}^-$ and bremsstrahlung cooling $dQ_{br}^- + dQ_{br,C}^-$. The dQ^- 's are in ergs s^{-1} . The mass accretion rate and the position of star r_* are shown in each panel.

Fig. 2.— We show that the electron temperatures in the left column and the radio spectra of the ADAFs in the right column in ergs s^{-1} in log scales. The continuous line and the dotted line indicate the electron temperature and the radio spectrum of the ADAFs with the stellar cooling, and of the ADAFs without the stellar cooling, respectively. Note that the difference between the two accretion flows with and without the stellar cooling is indistinguishable when $\dot{m} = 10^{-3}$.



